

Coherent states measurement entropy

Jarosław Kwapien^{||}, Wojciech Słomczyński[¶] and Karol Życzkowski[§]+

[†] Instytut Fizyki Jądrowej im. H Niewodniczańskiego, ul. Radzikowskiego 152, PL-31 305, Kraków, Poland

[‡] Instytut Matematyki, Uniwersytet Jagielloński, ul. Reymonta 4, PL-30 057, Kraków, Poland

[§] Instytut Fizyki im. M Smoluchowskiego, Uniwersytet Jagielloński, ul. Reymonta 4, PL-30 057, Kraków, Poland

Received 23 April 1996

Abstract. Coherent states (CS) quantum entropy can be split into two components. The dynamical entropy is linked with the dynamical properties of a quantum system. The measurement entropy, which tends to zero in the semiclassical limit, describes the unpredictability induced by the process of a quantum approximate measurement. We study the CS-measurement entropy for spin coherent states defined on the sphere discussing different methods dealing with the limit: time tends to infinity. In particular, we propose an effective technique of computing the entropy by iterated function systems. The dependence of CS-measurement entropy on the character of the partition of the phase space is analysed.

1. Introduction

During the last decade a lot of attention has been paid to the analysis of quantum analogues of chaotic classical maps defined on a compact phase space. In particular, quantum versions of the Baker map [1–3], the Arnold cat map [4, 3] (torus) and the periodically kicked top [5–7] (sphere) become standard models often used in the study on *quantum chaology* [8–10]. The classical versions of these models can be called *chaotic* (for the kicked top under an appropriate choice of parameters), since the Kolmogorov–Sinai (KS) dynamical entropy of the systems is positive. The definition of KS-entropy cannot be adopted straightforwardly into quantum mechanics, as it is based on the concept of classical trajectory. Several methods of generalizing KS-entropy to quantum mechanics were proposed (see [11–15] for the complete bibliography), but most of them lead to zero entropy for all quantum systems represented on a finite-dimensional Hilbert space. This property is common to the definitions of quantum entropy due to Connes, Narnhoffer and Thirring [16], Gaspard [17], Alicki and Fannes [18], and Roepstorff [19]. Therefore these concepts of quantum entropy are not suitable to describe dynamical properties of the above-mentioned quantum maps which, due to the compactness of the classical phase space, act on a finite-dimensional Hilbert space.

Coherent states (CS) dynamical entropy, introduced in [13], appears to be more adequate for a quantitative characterization of chaos in such quantum systems. This definition of quantum entropy takes into account the process of *sequential approximate measurement*. The notion of approximate, or unsharp, or fuzzy quantum measurement has been analysed

^{||} E-mail address: kwapien@castor.if.uj.edu.pl

[¶] E-mail address: slomczyn@im.uj.edu.pl

⁺ E-mail address: karol@castor.if.uj.edu.pl

in the last 25 yr by Ali, Emchs and Prugovečki, Busch and Lahti, Davies and Lewis, Ozawa, Schroeck and many others in the framework of the operational approach to quantum mechanics, or stochastic (phase space) quantum mechanics (see monographs [20–23] for details and further references). In order to get some information about the localization of a state in a phase space one may perform double (or multiple, in a phase space of higher dimension) quantum measurement of canonically conjugated observables. Due to the uncertainty principle such a measurement cannot be sharp and has to be approximate. The definition of CS-entropy is therefore based on the modified postulate of wavefunction collapse. The original postulate of Lüders and von Neumann, corresponding to a single exact measurement, assumes that after a measurement the state undergoes a transition to an eigenstate of the observable [24]. The modified postulate, used in the description of an approximate multiple measurement, asserts that after the measurement the state is transformed into an appropriate mixture of coherent states, i.e. the coherent states are *a posteriori* states in the sense of Ozawa [25].

The CS-entropy with respect to a given partition can be divided into two parts: the *dynamical entropy* which describes the dynamical properties of a quantum system and the *measurement entropy* related to the unpredictability induced by the process of sequential quantum measurement. The proof of the fact that in the semiclassical limit the CS-measurement entropy tends to zero was sketched in [13]. In the same paper it was conjectured that *the CS-dynamical entropy tends to the KS-entropy of the corresponding classical system* if the unitary dynamics comes from an appropriate quantization procedure, and some results in this direction were obtained.

In this work we analyse in detail CS-measurement entropy for the $SU(2)$ coherent states, where the phase space is the two-dimensional sphere. Such an example is of special physical interest, since it corresponds to an unsharp measurement of the spin components [26–29]. In [13] we proposed a general plan for studying the notion of CS-entropy. The results obtained here constitute the first step towards the realization of this scheme.

This paper is organized as follows. In section 2 we recall the definitions of CS-entropy, CS-measurement entropy and CS-dynamical entropy, and review some of their basic properties. In section 3 we summarize the standard facts on the $SU(2)$ vector coherent states. Various methods of computing CS-measurement entropy for the partition of the sphere into two hemispheres are presented, and the semiclassical limit is discussed in section 4. The case of an arbitrary partition of the sphere is analysed in section 5. In section 6 we treat yet another method of calculating CS-entropy based on the notion of iterated function systems. The Rényi-type generalizations of CS-entropy are introduced in section 7. Finally, section 8 contains some concluding remarks.

2. Coherent states quantum entropy

2.1. Kolmogorov–Sinai entropy

Coherent states entropy can be regarded [13] as a generalization of the classical Kolmogorov–Sinai entropy. Let us recall here the definition of the KS-entropy for a classical measurable map $S : \Omega \rightarrow \Omega$ generating a discrete dynamical system. Let Ω be a compact phase space endowed with a probability measure μ , invariant with respect to S , and divided into k disjoint measurable cells E_1, \dots, E_k . The time evolution of classical trajectories during n periods is described via probabilities

$$P_{i_0, \dots, i_{n-1}}^{cl} = \mu(\{x \in \Omega : x \in E_{i_0}, S(x) \in E_{i_1}, \dots, S^{n-1}(x) \in E_{i_{n-1}}\}) \quad (2.1)$$

of entering a given sequence of cells, where $i_l = 1, \dots, k; l = 0, \dots, n - 1$. It is assumed that the initial points, determining uniquely each trajectory, are distributed uniformly in the phase space with respect to the measure μ .

The partial entropy of S is

$$H_n = - \sum_{i_0, \dots, i_{n-1}=1}^k P_{i_0, \dots, i_{n-1}}^{cl} \ln P_{i_0, \dots, i_{n-1}}^{cl} \tag{2.2}$$

the *KS-entropy with respect to the partition* $\mathcal{C} = \{E_1, \dots, E_k\}$ is given by

$$H_{KS}(S, \mathcal{C}) := \lim_{n \rightarrow \infty} \frac{1}{n} H_n \tag{2.3}$$

and finally the *KS-entropy* of S is defined as [30]

$$H_{KS}(S) := \sup_{\mathcal{C}} H_{KS}(S, \mathcal{C}). \tag{2.4}$$

In the above formula the supremum is taken over all possible finite partitions of the phase space. A partition for which the supremum is achieved is called *generating*. Knowledge of a k -element generating partition for a given map allows one to represent the time evolution of the system in a k -letters symbolic dynamics and to find the upper bound for the KS-entropy: $H_{KS}(S) \leq \ln k$. For some classical systems, like the Baker map, it is straightforward to find a generating partition and to compute the KS-entropy. On the other hand, it is usually difficult to find a generating partition for an arbitrary classical map. Recent years brought some progress in this field: Christiansen and Politi found a good approximation for a generating partition for the standard map [31] and obtained a fair estimate for the KS-entropy (see also [32]).

The convergence to the limit in (2.3) is usually slow, not faster than $1/n$. It is therefore advantageous to consider the relative entropies G_n defined as

$$G_n := H_n - H_{n-1} \quad \text{for } n > 1 \quad G_1 = H_1. \tag{2.5}$$

It is easy to show that the sequence G_n also tends to H_{KS} [33]. This limit is usually achieved much faster than the limit in (2.3). For example Misiurewicz and Ziemian [34] and Ziemian [35] proved that for a certain class of maps from the unit interval onto itself this convergence is exponential (see also [36]). It seems that such a behaviour is typical for chaotic maps. We refer the reader to [37–40] for the review of recent results in this area. Note that the convergence of $\frac{1}{n} H_n = \frac{1}{n} \sum_{i=1}^n G_i$ is slower, since the terms of larger i have to balance a poor precision of the approximation due to the initial terms [41].

2.2. Approximate measurement and coherent states

The probabilities P^{cl} entering the definition of classical KS-entropy (2.2) are meaningful under the assumption that during the time evolution of the system one can trace an individual trajectory and determine its localization in the phase space with infinite precision. This supposition, consistent with the principles of classical mechanics, is definitely not fulfilled in quantum mechanics.

Information concerning the time evolution of a quantum system may be obtained by the process of sequential measurement. The fundamental analysis of a single quantum measurement of a discrete observable \hat{A} , expanded in an orthonormal basis as $\hat{A} := \sum_{m=1}^N a_m |m\rangle \langle m|$, leads to the *collapse postulate* of Lüders and von Neumann. The canonical measurement of \hat{A} yields with the probability $p_a = \sum_{a_m=a} \langle m | \hat{\rho} | m \rangle$ the state reduction [24]

$$\hat{\rho} \xrightarrow{\text{measurement}} \hat{\rho}' := \frac{\sum_{a_m=a} |m\rangle \langle m| \hat{\rho} |m\rangle \langle m|}{\sum_{a_m=a} \langle m | \hat{\rho} | m \rangle} \tag{2.6}$$

provided the outcome is a , where $\hat{\rho}$ is a density matrix describing the state of the system before the measurement. If $|m\rangle$ is the only eigenstate of \hat{A} corresponding to the eigenvalue a_m , then this formula simplifies and the act of measurement transforms $\hat{\rho}$ into a pure state $\hat{\rho}' = |m\rangle\langle m|$.

The measurement of a single observable does not provide sufficient information about the localization of the quantum state in the phase space. Such information can be acquired only in a simultaneous double (or multiple) approximate measurement of canonically conjugated observables. Let us consider an N -dimensional complex Hilbert space \mathcal{H} which represents the *kinematics* of the system and a compact set Ω equipped with a probability measure μ (we shall write dx for $d\mu(x)$) which forms a *phase space*, or in other words, a space of experimental outcomes. A correspondence between both spaces can be established by introducing a *family of coherent states*, i.e. a continuous map $\Omega \ni x \rightarrow |x\rangle \in \mathcal{H}$ satisfying the resolution of the identity $\int_{\Omega} |x\rangle\langle x| dx = I$ [42]. In this work we use the coherent states normalized as $\langle x|x\rangle = N$.

Following the ideas of Davies and Lewis [43] and Davies [20] we assume, in a full analogy to (2.6), that a multiple approximate quantum measurement yields the state reduction

$$\hat{\rho} \xrightarrow{\text{measurement}} \hat{\rho}' := \frac{\frac{1}{N} \int_{E_i} |x\rangle\langle x| \hat{\rho} |x\rangle\langle x| dx}{\int_{E_i} \langle x| \hat{\rho} |x\rangle dx} \quad (2.7)$$

provided the outcome is in the cell E_i , which occurs with the probability $P_i^{\text{CS}} = \int_{E_i} \langle x| \hat{\rho} |x\rangle dx$. Note that if one increases the precision of the measurement of a single variable (and simultaneously decreases the precision of the measurement of the canonically coupled variables) this postulate reduces in the limit to the standard collapse postulate of Lüders and von Neumann. Formally, one has to replace the coherent states $|x\rangle$ used in (2.7) by so-called *squeezed states*.

2.3. CS-probabilities and CS-entropy

Our approach to quantum entropy is based on the assumption that the knowledge about the time evolution of a quantum state is obtained from a sequence of multiple approximate quantum measurements. The evolution of the system between every two subsequent measurements is governed by a unitary matrix U .

A scheme of the first three periods of the time evolution of a system is presented in figure 1. Consider a quantum path encoded by the following sequence of cells: $\{E_{i_0}, E_{i_1}, \dots, E_{i_{n-1}}\}$. Let the initial state be proportional to the identity operator, i.e. $\hat{\rho}_0 =$




Figure 1. Scheme of the first three periods of the time evolution of the dynamical system. The unitary quantum map U describes the evolution of the system during each period, after which an act of approximate measurement takes place.

$1/N \cdot I$. The coherent states collapse postulate (2.7) allows us to calculate the probability that a given sequence of n symbols occurs ([13], see also [44]). Namely, we have

$$P_{i_0, \dots, i_{n-1}}^{\text{CS}} = \int_{E_{i_0}} dx_0 \cdots \int_{E_{i_{n-1}}} dx_{n-1} \prod_{u=1}^{n-1} K(x_{u-1}, x_u) \tag{2.8}$$

where $i_l = 1, \dots, k; l = 0, \dots, n - 1$ and the kernel K is given by

$$K(x, y) = \frac{1}{N} |\langle y|U|x \rangle|^2. \tag{2.9}$$

We call them *CS-probabilities*. Partition-dependent, *coherent states (CS) entropy* H^{CS} of a quantum map U is defined like its classical counterpart (2.3)

$$H^{\text{CS}}(U, \mathcal{C}) := \lim_{n \rightarrow \infty} \frac{1}{n} H_n(U, \mathcal{C}) \tag{2.10}$$

where

$$H_n(U, \mathcal{C}) := - \sum_{i_0, \dots, i_{n-1}=1}^k P_{i_0, \dots, i_{n-1}}^{\text{CS}} \ln P_{i_0, \dots, i_{n-1}}^{\text{CS}} \tag{2.11}$$

and $\mathcal{C} = \{E_1, \dots, E_k\}$. In the semiclassical limit the CS-entropy seems to tend to the KS-entropy, if the quantization procedure is *regular* [13], i.e. if some assumptions linking the family of quantum maps with the corresponding classical map are fulfilled.

Quantum CS-probabilities can be also used to define other quantities which measure the randomness of the system (for a recent account of such concepts see [45,46]) like Rényi-type entropy of order β which we shall analyse in section 7. For some purposes, for instance, it might be useful to define *CS-inverse participation ratio* ν

$$\nu(U, \mathcal{C}) := \sum_{i_0, \dots, i_{n-1}=1}^k (P_{i_0, \dots, i_{n-1}}^{\text{CS}})^2. \tag{2.12}$$

It is an analogue of a quantity often used in solid state physics to describe localization of a wavefunction [47], since its inverse gives the average number of occupied cells. It is linked to CS-Rényi entropy of order 2.

In the simplest case of the trivial dynamics the quantum map U reduces to the identity operator I . Even in this case the quantum entropy H^{CS} does not vanish, since the coherent states are not orthogonal and do overlap [42]. The *CS-measurement entropy* is given by [13,48]

$$H_{\text{meas}}(\mathcal{C}) := H^{\text{CS}}(U \equiv I, \mathcal{C}) \tag{2.13}$$

and depends on a family of coherent states in the phase space Ω and on a finite partition \mathcal{C} .

The *CS-dynamical entropy of a quantum map U with respect to a partition \mathcal{C}* is defined as [13,48]

$$H_{\text{dyn}}(U, \mathcal{C}) := H^{\text{CS}}(U, \mathcal{C}) - H_{\text{meas}}(\mathcal{C}) \tag{2.14}$$

and partition-independent *CS-dynamical entropy* as [49]

$$H_{\text{dyn}}(U) := \sup_{\mathcal{C}} H_{\text{dyn}}(U, \mathcal{C}). \tag{2.15}$$

In the present paper we study CS-measurement entropy and its dependence on a partition and the semiclassical parameter. This is a preliminary step to calculating CS-dynamical entropy, which is defined as the difference of two quantities. Moreover, the techniques we use in computing of CS-measurement entropy can also be applied in the general case.

2.4. Properties of CS-measurement entropy

We now review some basic properties of CS-measurement entropy. Let us assume that a finite partition \mathcal{C} of the phase space Ω is given. Let $H_n(\mathcal{C})$ be defined by (2.8)–(2.11) with $U = I$, and let $G_1(\mathcal{C}) = H_1(\mathcal{C})$; $G_n(\mathcal{C}) = H_n(\mathcal{C}) - H_{n-1}(\mathcal{C})$, for $n > 1$. Then, applying the general theory of entropy for random transformations [50, 51], we obtain the following facts:

- (1) the sequences $\frac{1}{n}H_n(\mathcal{C})$ and $G_n(\mathcal{C})$ decrease with n to $H_{\text{meas}}(\mathcal{C})$;
- (2) if a partition \mathcal{C}' is finer than a partition \mathcal{C} , then $H_{\text{meas}}(\mathcal{C}') \geq H_{\text{meas}}(\mathcal{C})$.

Next, let us observe that the kernel K which appears in (2.8) is bistochastic, i.e. $\int_{\Omega} K(x, \bar{y}) dx = \int_{\Omega} K(\bar{x}, y) dy = 1$ for all $\bar{x}, \bar{y} \in \Omega$. Let us denote by K_0 the maximum of K . Then

- (3) the CS-measurement entropy fulfills the following inequalities:

$$\frac{1}{n}H_n(\mathcal{C}) - \frac{1}{n} \ln K_0 \leq H_{\text{meas}}(\mathcal{C}) \leq \frac{1}{n}H_n(\mathcal{C}) \quad (2.16)$$

and, in consequence,

$$H_1(\mathcal{C}) - \ln K_0 \leq H_{\text{meas}}(\mathcal{C}) \leq H_1(\mathcal{C}) \quad (2.17)$$

(for the proof see appendix A).

Note that $H_1(\mathcal{C})$ does not depend on the family of coherent states but only on the measure μ and it is just the entropy of the partition \mathcal{C} with respect to the measure μ . If Ω is a Riemannian manifold and μ is the Riemannian measure on Ω , then one can deduce from (2) and (3) that $H_{\text{meas}}(\mathcal{C})$ can be arbitrarily large for a sufficiently fine partition \mathcal{C} .

It follows from (1) and (3) that

- (4) if $H_1(\mathcal{C}) \neq H_{\text{meas}}(\mathcal{C})$, then the sequence $\frac{1}{n}H_n(\mathcal{C})$ converges to the entropy $H_{\text{meas}}(\mathcal{C})$ precisely as n^{-1} .

(For the proof see appendix B.)

Another important property of CS-measurement entropy,

- (5) $H_{\text{meas}}(\mathcal{C})$ tends to 0 in the semiclassical limit,

was proved in [13] for $SU(2)$ (spin) coherent states. The decay seems to be rather slow. We shall try to evaluate its rate in the following.

2.5. Matrix form of CS-probabilities

Let $\mathcal{C} = \{E_1, \dots, E_k\}$. We assume that the kernel K entering formula (2.8) has the form

$$K(x, y) = \sum_{l,r=0}^M a_{lr} g_r(x) f_l(y) \quad \text{for } x, y \in X \quad (2.18)$$

where $a_{lr} \in \mathbb{R}$, $f_l, g_r : \Omega \rightarrow \mathbb{R}$ are continuous, for $l, r = 0, \dots, M$, and $f_0 = g_0 \equiv 1$ (in fact we can always present K in such a form if the family of coherent states comes from the canonical group-theoretic construction (see [52–54]) with a finite-dimensional Hilbert space \mathcal{H} ; then M is an increasing function of the dimension of the Hilbert space). Let us define matrices $A = [a_{lr}]_{l,r=0}^M$ and $B(i)_{rl} = \int_{E_i} g_r(x) f_l(x) dx$ for $l, r = 0, \dots, M$, $i = 1, \dots, k$. Then the CS-probabilities are given by the first element of the following matrix product:

$$P_{i_0, \dots, i_{n-1}}^{\text{CS}} = (B(i_{n-1})AB(i_{n-2})A \dots AB(i_0))_{00}. \quad (2.19)$$

(The proof will appear in [55].) Now one can show that the family of the CS-probabilities generates on the code space $\mathcal{S}^{\mathbb{N}}$, where $\mathcal{S} = \{1, \dots, k\}$, a shift-invariant measure, which is *algebraic* in the sense of Fannes *et al* (see [56]). Clearly, the decomposition of the kernel

K is not unique. Moreover, the assumption $f_0 = g_0 \equiv 1$ is too restrictive. In fact, to apply the matrix method, it is enough to know that the constant function 1 is a linear combination of the functions f_0, \dots, f_M [55].

The above formula makes the calculation of the CS-entropy much easier. Moreover, it is a starting point for the further investigation of entropy utilizing the theory of iterated function systems. We present in section 6 some results in this direction. For a fuller treatment we refer the reader to [55].

3. Spin coherent states

The two-dimensional sphere S^2 can be considered as the phase space of the periodically kicked top. This classical dynamical system is known to exhibit chaos under a suitable choice of system parameters [5]. In order to study a quantum analogue of this system it is convenient to consider the operator of angular momentum J . Its three components $\{J_x, J_y, J_z\}$ are related to the infinitesimal rotations along three orthogonal axes $\{x, y, z\}$ in \mathbb{R}^3 and fulfil the standard commutation relations $[J_l, J_m] = i\epsilon_{lmn}J_n$, where $l, m, n = x, y, z$ and ϵ_{lmn} represents the antisymmetric tensor (from now on we put $\hbar = 1$). The operators $J_{\pm} = J_x \pm iJ_y$ and J_z are generators of the compact Lie group $SU(2)$. The eigenvalues $j(j+1)$, $j = 0, \frac{1}{2}, 1, \frac{3}{2}, \dots$, of the Casimir operator $J^2 = J_x^2 + J_y^2 + J_z^2$ determine the dimension $N = 2j + 1$ of the Hilbert spaces \mathcal{H}_N carrying the representation of the group. Common eigenstates $|j, m\rangle$, $m = -j, \dots, j$, of the operators J^2 and J_z form an orthonormal basis in \mathcal{H}_N .

The $SU(2)$ (spin) coherent states were introduced by Radcliffe [57] and Arecchi *et al* [58]. For a thorough discussion we refer the reader to [42, 52–54, 59, 60]. The idea is the following. Each point on the sphere labelled by the spherical coordinates (ϑ, φ) corresponds to the $SU(2)$ coherent state $|j, \vartheta, \varphi\rangle$ generated by the unitary operator $R(\vartheta, \varphi) = \exp[i\vartheta(\sin\varphi J_x - \cos\varphi J_y)]$ acting on the reference state $|j, j\rangle$. The natural projection $SU(2) \rightarrow SO(3)$ relates with the operator $R(\vartheta, \varphi)$ the rotation by the angle ϑ around the axis directed along the vector $(\sin\varphi, -\cos\varphi, 0)$ normal to the z -axis and to the vector $(\sin\vartheta\cos\varphi, \sin\vartheta\sin\varphi, \cos\vartheta)$ (see figure 2). The state $|j, j\rangle$, pointing towards the ‘north pole’ of the sphere, enjoys the minimal uncertainty, i.e. the expression $\sum_{l=x,y,z} \Delta J_l^2$ takes in this state the minimal value j (the other possible choice of the reference state is $|j, -j\rangle$). More precisely, we put

$$|j, \vartheta, \varphi\rangle = \sqrt{2j+1}R(\vartheta, \varphi)|j, j\rangle. \tag{3.1}$$

Using the stereographical projection $\gamma = \tan(\vartheta/2)\exp(i\varphi)$ one can find a complex representation of the coherent state $|j, \gamma\rangle := |j, \vartheta, \varphi\rangle$

$$|j, \gamma\rangle = \frac{\sqrt{2j+1}}{(1+|\gamma|^2)^j} \exp[\gamma J_-]|j, j\rangle. \tag{3.2}$$

The prefactor $\sqrt{2j+1}$ introduced into the above formulae ensures the coherent states identity resolution in the form

$$\int_{S^2} |j, \vartheta, \varphi\rangle\langle j, \vartheta, \varphi| d\mu(\vartheta, \varphi) = I \tag{3.3}$$

where the Riemannian measure μ on S^2 is given by $d\mu = \sin\vartheta d\vartheta d\varphi/4\pi$ and therefore does not depend on the quantum number j . The norm of the coherent states changes with j as $|\langle j, \vartheta, \varphi|j, \vartheta, \varphi\rangle| = 2j+1$, which enables the respective *Husimi-like distribution* $S^2 \ni (\vartheta', \varphi') \rightarrow |\langle j, \vartheta, \varphi|j, \vartheta', \varphi'\rangle|^2 \in \mathbb{R}$ of the coherent state $|j, \vartheta, \varphi\rangle$ to tend to the




Figure 2. Spherical representation of the spin coherent state $|\vartheta, \varphi\rangle$ generated by the unitary rotation operator $R(\vartheta, \varphi)$.

Dirac δ -function as $j \rightarrow \infty$. Thus, after such a renormalization we can treat the limit $j \rightarrow \infty$ as the *semiclassical limit* [54] or, in other words, as the *sharp-point limit* in the sense of Schroeck [61]. For an interpretation of this limit in the language of nonstandard analysis see [62]. If we had transformed spin coherent states in a different way defining $||j, \sqrt{2j}\gamma\rangle := (1+|\gamma|^2)^{-j} \exp[\gamma J_-]|j, j\rangle$, we would have obtained the canonical (harmonic oscillator) coherent states in the limit $j \rightarrow \infty$. This kind of limit, however, is completely different from the semiclassical limit we use in the present paper.

To simplify the notation in the following sections we shall omit the number j labelling coherent states $|\vartheta, \varphi\rangle$ or $|\gamma\rangle$. Note that S^2 is isomorphic to the coset space $SU(2)/U(1)$, where $U(1)$ is the maximal stability subgroup of $SU(2)$ with respect to the state $|j, j\rangle$, i.e. the subgroup of all elements of $SU(2)$ which leave $|j, j\rangle$ invariant up to a phase factor. Hence the above construction can be treated as a particular case of the general construction of group-theoretic coherent states.

Expansion of a coherent state in the eigenbasis of J^2 and J_z reads

$$|\vartheta, \varphi\rangle = \sqrt{2j+1} \sum_{m=-j}^{m=j} \sin^{j-m} \left(\frac{\vartheta}{2}\right) \cos^{j+m} \left(\frac{\vartheta}{2}\right) \exp(i(j-m)\varphi) \left[\binom{2j}{j-m} \right]^{1/2} |j, m\rangle. \quad (3.4)$$

The expectation values of the components of J are

$$\langle j, \vartheta, \varphi | J | j, \vartheta, \varphi \rangle = j(2j+1)(\sin \vartheta \cos \varphi, \sin \vartheta \sin \varphi, \cos \vartheta) \quad (3.5)$$

which establishes the link between the coherent state $|j, \vartheta, \varphi\rangle$ and the vector (ϑ, φ) oriented along the direction defined by a point on the sphere.

The infinite basis formed in the Hilbert space by the coherent states is overcomplete. Two different $SU(2)$ coherent states overlap unless they point towards two opposite poles

on the sphere. Expanding two coherent states in the $|j, m\rangle$ basis (3.4) we can calculate their overlap as

$$|\langle \vartheta', \varphi' | \vartheta, \varphi \rangle|^2 = (2j + 1)^2 \left(\frac{1 + \cos \Xi}{2} \right)^{2j} \tag{3.6}$$

where Ξ is the angle between two vectors on S^2 related to the coherent states $|\vartheta, \varphi\rangle$ and $|\vartheta', \varphi'\rangle$. Hence the transition kernel K defined by (2.9) takes (for $U = I$) the form

$$\begin{aligned} K((\vartheta, \varphi), (\vartheta', \varphi')) &= \frac{|\langle \vartheta', \varphi' | \vartheta, \varphi \rangle|^2}{2j + 1} \\ &= \frac{2j + 1}{2^{2j}} [1 + \cos \vartheta \cos \vartheta' + \sin \vartheta \sin \vartheta' \cos(\varphi - \varphi')]^{2j}. \end{aligned} \tag{3.7}$$

The overlap decreases to 0 with j for $|\vartheta, \varphi\rangle \neq |\vartheta', \varphi'\rangle$ and sufficiently large j .

4. Measurement entropy for two hemispheres

We would like to compute the CS-measurement entropy for the case corresponding to the physical process of simultaneous approximate measurement of different spin components. Let us first consider the simplest case, where the classical phase space Ω equal to the two-dimensional sphere S^2 is divided into two hemispheres $E_+ = \{(\vartheta, \varphi) : \varphi \in [0, 2\pi), \vartheta \in [0, \pi/2]\}$ and $E_- = \{(\vartheta, \varphi) : \varphi \in [0, 2\pi), \vartheta \in (\pi/2, \pi]\}$. The result of any measurement $i = \pm 1$ gives information about the orientation of the spin.

4.1. Transition probabilities

The CS-transition probabilities P^{CS} for the results i_0, \dots, i_{n-1} of n consecutive measurements are obtained from (2.8) and (2.9) by setting the evolution operator U to be the identity and taking the appropriate integration domains. The explicit integral reads

$$\begin{aligned} P_{i_0, \dots, i_{n-1}}^{CS} &= (4\pi)^{-n} \int_{E_{i_0}} \sin \vartheta_0 \, d\vartheta_0 \, d\varphi_0 \dots \\ &\dots \int_{E_{i_{n-1}}} \sin \vartheta_{n-1} \, d\vartheta_{n-1} \, d\varphi_{n-1} \prod_{u=1}^{n-1} K((\vartheta_{u-1}, \varphi_{u-1}), (\vartheta_u, \varphi_u)) \end{aligned} \tag{4.1}$$

where the kernel K is given by (3.7), and $i_u = \pm 1$ for $u = 0, \dots, n - 1$. Straightforward integration allows one to obtain analytical results for low values of n and j [48].

In spite of the trivial dynamics ($U \equiv I$) the result of the first measurement may differ from the second one, and consequently, all the transition probabilities are nonzero. In the semiclassical limit $j \rightarrow \infty$ the ‘mixed’ transition probabilities (e.g. $P_{+-}^{CS} = P_{-+}^{CS}$) vanish, while the survival probabilities (e.g. $P_{++}^{CS}, P_{+++}^{CS}$) tend to $\frac{1}{2}$. The geometric symmetry of reflection induces the invariance of the probabilities with respect to the interchange of signs ($+ \longleftrightarrow -$). Moreover, due to the time-reversal invariance, the CS-probability for any sequence of results equals the CS-probability of the same sequence written in the reverse order (e.g. $P_{++++}^{CS} = P_{+---}^{CS}$). One can observe that for a given number of measurements n , the probabilities for two sequences of results with the same number of transitions are similar (e.g. for one transition: $P_{+---}^{CS} \approx P_{+++}^{CS}$; for two transitions: $P_{+---+}^{CS} \approx P_{++++}^{CS}$). Direct integration of (4.1) does not allow one to obtain the CS-probabilities for larger values of j or n , which is necessary to estimate the CS-measurement entropy. For this purpose it is convenient to formulate integrals in matrix form.

4.2. Matrix formulation of integrals

Computation of the CS-probabilities can be significantly simplified by applying the general method described in section 2.5. This can be seen, especially, for the division of the sphere into several latitudinal components E_1, \dots, E_k , where $E_i = \{(\vartheta, \varphi) : \varphi \in [0, 2\pi), \vartheta \in W_i\}$ for $i = 1, \dots, k$, and $\{W_1, \dots, W_k\}$ is a partition of the interval $[0, \pi]$. Performing the substitutions $t_i = \cos \vartheta_i$ and integrating over $\varphi_0, \dots, \varphi_n$ we can simplify formula (4.1) writing

$$P_{i_0, \dots, i_{n-1}}^{\text{CS}} = \int_{\tilde{W}_{i_0}} \frac{1}{2} dt_0 \dots \int_{\tilde{W}_{i_{n-1}}} \frac{1}{2} dt_{n-1} \prod_{u=1}^{n-1} \tilde{K}(t_{u-1}, t_u) \quad (4.2)$$

where $\tilde{W}_i = \{\cos t : t \in W_i\}$ for $i = 1, \dots, k$, and the reduced kernel \tilde{K} is given by

$$\tilde{K}(t, s) = \frac{2j+1}{4^{2j}} \sum_{q=0}^{2j} \binom{2j}{q}^2 ((1+t)(1+s))^q ((1-t)(1-s))^{2j-q} = \sum_{l,r=0}^{2j} a_{lr} t^l s^r \quad (4.3)$$

for $t, s \in [-1, 1]$. Thus the kernel \tilde{K} is represented in the form (2.18) with $\tilde{\Omega} = [-1, 1]$, $d\tilde{\mu}(t) = \frac{1}{2} dt$, $f_l(t) = t^l$, $g_r(s) = s^r$ for $t, s \in \tilde{\Omega}$, and $M = 2j$. Note that $\{\tilde{W}_1, \dots, \tilde{W}_k\}$ forms a partition of $\tilde{\Omega}$. Hence we can apply formula (2.19) for the CS-probabilities writing them in the matrix form

$$P_{i_0, \dots, i_{n-1}}^{\text{CS}} = \langle (1, 0, \dots, 0) | (B(i_{n-1}) A B(i_{n-2}) A \dots A B(i_0)) | (1, 0, \dots, 0) \rangle \quad (4.4)$$

with $A = [a_{lr}]_{l,r=0}^{2j}$ (given by (4.3)) and $B(i)_{rl} = \frac{1}{2} \int_{\tilde{W}_i} t^{l+r} dt$ for $i = 1, \dots, k$; $l, r = 0, \dots, 2j$.

If we divide the sphere into two hemispheres, then $B(i)_{rl} = i^{l+r}/2(l+r+1)$ for $i = \pm 1$; $l, r = 0, \dots, 2j$. In this case (4.4) takes a particularly simple form for $j = \frac{1}{2}$

$$P_{i_0, \dots, i_{n-1}}^{\text{CS}} = \frac{1}{2^n} \langle (1, 0) | \begin{pmatrix} 1 & i_0/2 \\ i_0/2 & \frac{1}{3} \end{pmatrix} \begin{pmatrix} 1 & i_1/2 \\ i_1/2 & \frac{1}{3} \end{pmatrix} \dots \begin{pmatrix} 1 & i_{n-1}/2 \\ i_{n-1}/2 & \frac{1}{3} \end{pmatrix} | (1, 0) \rangle \quad (4.5)$$

where the results of the measurements i_u are equal to -1 or $+1$ for $u = 0, \dots, n-1$.

4.3. Limit $n \rightarrow \infty$

In the remainder of this section we assume that \mathcal{C} is the partition of the sphere into two hemispheres, i.e. $\mathcal{C} = \{E_+, E_-\}$. Moreover, we set $H_{\text{meas}} := H_{\text{meas}}(\mathcal{C})$, $H_n := H_n(\mathcal{C})$ and $G_n := G_n(\mathcal{C})$.

In table 1 we present partial and relative entropies calculated for two different values of j with the aid of the formulae (2.10), (2.11) and (4.4).

We assert in section 2.4 (4) that H_n converges to H_{meas} exactly as $1/n$. One can deduce from table 1 that the convergence of G_n to the same limit is much faster. In fact, it seems to be exponential. In section 6 we give some arguments supporting this statement. Thus, to calculate the limiting value we use the extrapolations $H_n \sim H_{\text{meas}} + \alpha/n$ and $G_n \sim H_{\text{meas}} + \gamma c^n$. The outcomes are contained in table 1. Let us observe that the rate of convergence decreases with j and hence the method of computing the CS-measurement entropy based on formula (4.4) does not lead to satisfactory results in the semiclassical limit, i.e. for large quantum number j .

Table 1. Partial entropy H_n/n and relative entropy G_n for the partition $\mathcal{C} = \{E_+, E_-\}$, the quantum number $j = \frac{1}{2}$ and $j = 10$, and the number of measurements $n = 1, \dots, 8$ with an extrapolation to $n \rightarrow \infty$.

n	$j = \frac{1}{2}$		$j = 10$	
	H_n/n	G_n	H_n/n	G_n
1	0.693 147 180 559	0.693 147 180 559	0.693 1471	0.693 1471
2	0.677 355 209 358	0.661 563 238 157	0.532 3993	0.371 6514
3	0.672 009 066 259	0.661 316 780 060	0.473 4456	0.355 5383
4	0.669 335 388 698	0.661 314 356 017	0.442 9253	0.351 3642
5	0.667 731 177 545	0.661 314 332 934	0.424 3127	0.349 8623
6	0.666 661 703 407	0.661 314 332 713	0.411 8012	0.349 2438
7	0.665 897 793 307	0.661 314 332 711	0.402 8258	0.348 9737
8	0.665 324 860 733	0.661 314 332 711	0.396 0793	0.348 8532
\vdots	\vdots	\vdots	\vdots	\vdots
∞	0.661 3	0.661 314 332 711	0.348 8	0.348 7560

4.4. Semiclassical regime $j \gg 1$

The matrix formula for the CS-probabilities is useful in numerical calculations, but as was mentioned above, does not allow us to compute the entropy for very large values of j . For two measurements, however, one can obtain some exact results. Applying (4.2) and (4.3) we get an analytical formula for the CS-probability valid for any j :

$$P_{+-}^{\text{CS}} = \binom{4j+1}{2j} 2^{-4j-2}. \tag{4.6}$$

(For the proof see appendix C.)

Due to symmetry $P_{-+}^{\text{CS}} = P_{+-}^{\text{CS}}$ and $P_{++}^{\text{CS}} = P_{--}^{\text{CS}} = \frac{1}{2} - P_{+-}^{\text{CS}}$. It is convenient to introduce a j -dependent coefficient $\tau_j = P_{+-}^{\text{CS}}/P_{++}^{\text{CS}}$, which tends to zero in the semiclassical limit $j \rightarrow \infty$. Using formula (4.6) we obtain

$$\tau_j = \frac{\binom{4j+1}{2j}}{2^{4j+1} - \binom{4j+1}{2j}}. \tag{4.7}$$

In order to get an upper bound for the CS-measurement entropy we may compute the relative entropy $G_2 = H_2 - H_1$ (see section 2.4). The partial entropy after one measurement H_1 equals $\ln 2$, independently of j . Summing over four possible paths $++$, $+-$, $-+$, $--$ one can compute the partial entropy H_2 obtaining finally the formula

$$G_2 = \ln(\tau_j + 1) - \frac{\tau_j}{\tau_j + 1} \ln(\tau_j) \tag{4.8}$$

which is symmetric with respect to an involution $\tau_j \rightarrow 1/\tau_j$.

Inserting the expression (4.7) into the above formula we get an explicit approximation for H_{meas} . It is represented by a full curve in figure 3, while circles denote the results obtained numerically for small j with the help of the matrix method presented above. In the semiclassical range $j \gg 1$ it is legitimate to apply the Stirling approximation of the factorial in (4.7), which gives a fair approximation and an upper bound for the CS-measurement entropy

$$H_{\text{meas}} < G_2 \sim \frac{\ln j}{2\sqrt{2\pi j}}. \tag{4.9}$$




Figure 3. CS-measurement entropy H_{meas} for two hemispheres as a function of the quantum number j . Circles represent numerical results, while the full curve stands for an upper bound G_2 given by (4.8).

It is worth noting that formula (4.8) can be obtained from the Markovian approximation of the CS-probabilities. Let us assume for a moment that the probabilities $P_{i_0, \dots, i_{n-1}}^{\text{CS}}$ were generated by a Markov shift. It follows from the symmetry of the problem that its initial vector would be $(\frac{1}{2}, \frac{1}{2})$ and its transition matrix Q would have the form

$$Q = \begin{pmatrix} a & 1-a \\ 1-a & a \end{pmatrix} \quad \text{where } a = 2P_{++}^{\text{CS}} = \frac{1}{\tau_j + 1}. \quad (4.10)$$

In fact, our probabilities $P_{i_0, \dots, i_{n-1}}^{\text{CS}}$ are not generated by a Markov shift; nevertheless, one can consider the Markovian approximation as above. Then the approximate probabilities P^{Mar} depending only on the number L of ‘transitions’ from one hemisphere to the other ($L = \frac{1}{2}(n-1 - \sum_{q=1}^{n-1} i_q i_{q-1})$) are equal to

$$P_{i_0, \dots, i_{n-1}}^{\text{Mar}} = \frac{1}{2} Q_{i_0, i_1} \times \dots \times Q_{i_{n-2}, i_{n-1}} = \frac{1}{2} (1-a)^L a^{n-1-L} = \frac{\tau_j^L}{2(\tau_j + 1)^{n-1}}. \quad (4.11)$$

In this approximation the probabilities form a geometric series with the same ratio τ_j for any number of measurements n .

Summing over all 2^n possible sequences we obtain the following approximate formula for the partial entropy:

$$\begin{aligned} H_n^{\text{Mar}} &= - \sum_{L=0}^{n-1} \binom{n-1}{L} \frac{\tau_j^L}{(\tau_j + 1)^{n-1}} \ln \left[\frac{\tau_j^L}{2(\tau_j + 1)^{n-1}} \right] \\ &= \ln 2 + (n-1) \left[\ln(\tau_j + 1) - \frac{\tau_j}{\tau_j + 1} \ln(\tau_j) \right]. \end{aligned} \quad (4.12)$$

Now dividing both sides by n and performing the limit $n \rightarrow \infty$ we arrive at the relative entropy G_2 given by (4.8).

Let us recall that in the semiclassical limit ($j \rightarrow \infty$) the relative measurement entropy G_2 tends to zero as $\ln j / \sqrt{j}$. This defines the scale in which the quantum effects are revealed. Unfortunately the precision of this approximation is not sufficient to conclude whether or not the logarithmic prefactor describes correctly the decay of the measurement entropy H_{meas} in the semiclassical limit, or whether its existence is an artifact introduced by the approximation.




Figure 4. Partition of the sphere divided along a parallel Θ_c into two connected cells.




Figure 5. Partial entropy H_n/n for the partition presented in figure 4 as a function of the parameter $\cos \Theta_c$ ($n = 8$). The values of j are given in the figure.

5. CS-measurement entropy for various partitions

The CS-measurement entropy depends on the number of cells in a partition and on their shape. In this section we consider several partitions of a different type. Figures 4 and 7 contain schemes for these partitions. In all the cases we compute the entropy using the matrix formulation introduced in sections 2.5 and 4.2. As in the preceding section we assume that \mathcal{C} denotes the respective partition of the sphere, putting $H_{\text{meas}} := H_{\text{meas}}(\mathcal{C})$, $H_n := H_n(\mathcal{C})$, and $G_n := G_n(\mathcal{C})$.

5.1. Two connected cells

Let us split the sphere into two segments along a parallel Θ_c . The northern segment E_+ contains points with $\vartheta \in [0, \Theta_c]$, while the southern E_- those with $\vartheta \in (\Theta_c, \pi]$. This partition is shown schematically in figure 4.

Figure 5 represents the dependence of the partial entropy H_n/n on the variable $\cos \Theta_c$ for $n = 8$ measurements and several values of j . For each value of n and j the partial entropy achieves its maximum at $\cos \Theta_c = 0$, for the partition into two hemispheres. The solid horizontal line drawn at $\ln 2$ represents the maximal entropy admissible for the partition containing two cells. For increasing values of j the partial entropy decreases and tends to




Figure 6. Partial entropies H_n/n and measurement entropy $H_{\text{meas}} = \lim_{n \rightarrow \infty} H_n/n$ for the partition presented in figure 4 with $j = \frac{1}{2}$ (light curves), and $j = 5$ (heavy curves).




Figure 7. Partition of the sphere into (a) two cells plotted out by parallels Θ_d and $\pi - \Theta_d$: a spherical zone and the union of two spherical segments; (b) two disconnected cells created by the equator and the spherical wedge of the radian measure Φ_d .

zero for $j \rightarrow \infty$.

For any of these partitions the partial entropy H_n/n approaches the limiting value H_{meas} approximately as $1/n$ (see section 2.4 (4)). As in the previously discussed case of two hemispheres, we estimate the limiting value H_{meas} by computing the relative entropy G_n . Figure 6 shows a comparison of the partial entropies H_n/n with H_{meas} extrapolated in this way for $j = \frac{1}{2}$ and $j = 5$. The difference increases with the spin length j .

5.2. Two disconnected cells

Let us now analyse another two classes of partitions of the sphere into two cells. In the first case (figure 7(a)) we divide the sphere into three parts along parallels $\pi - \Theta_d$ and Θ_d ,




Figure 8. CS-measurement entropy H_{meas} as a function of the number of cells k for $j = \frac{1}{2}$ (circles). The full curve represents the function $\ln k$. The inset shows the dependence of $H_{\text{max}} := \ln k - H_{\text{meas}}$ on k^{-2} obtained for $k = 30, \dots, 1000$.

and then join the lower and upper parts, thus obtaining two cells: a connected spherical zone and the disconnected union of two spherical segments. The CS-measurement entropy changes in this case from 0 (for $\Theta_d = 0$) to $\ln 2$ (for $\Theta_d = \pi/3$), which is the largest possible value for the CS-measurement entropy with respect to a two-element partition.

In the second case (figure 7(b)) we start from the splitting of the sphere into the lower and upper hemispheres. Next, we cut symmetrically two ‘pieces of cake’ out of both hemispheres, and then join the four parts across. We get in this way two disconnected cells marked in black and white in figure 7(b). The CS-measurement entropy changes from $\ln 2$ (for $\Phi_d = \pi$) to $0.6613\dots$ (for $\Phi_d = 0$). The latter case relates to the partition of the sphere into two hemispheres studied in section 4.

5.3. Many cells

Let us consider k disjoint zones created on the sphere by $k - 1$ parallels. As in the case of two cells, represented in figure 4, the CS-measurement entropy seems to achieve its maximum if the cells have the same volume $= 1/k$. We computed the CS-measurement entropy $H_{\text{meas}}(\mathcal{C}_k)$ for the partitions \mathcal{C}_k of the sphere into $k = 2, \dots, 1000$ zones of the same volume. Note that for large k even the second relative entropy $G_2(\mathcal{C}_k)$ provides a reliable estimate for $H_{\text{meas}}(\mathcal{C}_k)$. In figure 8 we present the CS-measurement entropy displayed for $j = \frac{1}{2}$ as a function of the number of cells k (circles). The full curve represents the function $\ln k$, which gives the upper bound for the entropy with respect to a partition consisting of k cells.

Since for any partition \mathcal{C} and a quantum map U the CS-dynamical entropy is defined by (2.14) as the difference of $H(U, \mathcal{C})$ and $H_{\text{meas}}(\mathcal{C})$, it is convenient to consider the quantity $H_{\text{max}}(\mathcal{C}_k) := \ln k - H_{\text{meas}}(\mathcal{C}_k)$, limiting the partition-dependent dynamical entropy $H_{\text{dyn}}(U, \mathcal{C}_k)$ from above. From (2.17) and (3.7) we know that $H_{\text{max}}(\mathcal{C}_k) \leq \ln(2j + 1)$ as $H_1(\mathcal{C}_k) = \ln k$. Although using this method one can establish the finiteness of the partition-independent dynamical entropy $H_{\text{dyn}}(U)$ given by (2.15), this upper bound seems to be rather crude. In fact $H_{\text{max}}(\mathcal{C}_k)$ decreases with j . It is interesting to observe that this quantity converges for $k \rightarrow \infty$. The limiting value depends on j and is close to 0.06 for $j = \frac{1}{2}$. The

inset in figure 8 shows the dependence of $H_{\max}(C_k)$ on k^{-2} for $k = 30, \dots, 1000$; $j = \frac{1}{2}$. The data displayed in this way are fitted well by a straight line, which allows us to postulate an approximate relation $H_{\text{meas}}(C_k) \approx \ln k - 0.05999745 + 0.1637/k^2$, found for $j = \frac{1}{2}$.

6. CS-entropy and iterated function systems

In this section we establish a relationship between CS-entropy and iterated function systems (IFSs). Firstly, we show how to obtain an IFS from a bistochastic kernel and a partition of the phase space. Then, we use this system to get an integral formula for CS-entropy and propose a new method of computing CS-entropy based on the ergodic theorem for IFSs. For more information on IFSs see [64–66].

6.1. Iterated function systems and an integral formula for CS-entropy

We follow the notation of sections 2.4 and 2.5. With each cell E_i ($i = 1, \dots, k$) of the partition we associate an $(M+1) \times (M+1)$ matrix $D(i) = B(i)A$. We consider functions $p_i : \mathbb{R}^M \rightarrow \mathbb{R}^+$ and partial maps $F_i : \mathbb{R}^M \rightarrow \mathbb{R}^M$ given by

$$p_i(\lambda) = (1, 0, \dots, 0)(D(i)(1, \lambda)) \quad (6.1)$$

and

$$F_i(\lambda) = (D(i)(1, \lambda))/p_i(\lambda) \quad (6.2)$$

for $\lambda \in \mathbb{R}^M$, $i = 1, \dots, k$.

Let us suppose that the functions $g_0 \equiv 1, g_1, \dots, g_M$ are linearly independent. Then one can show that

(0) $p_i(\lambda) \geq 0$ for $i = 1, \dots, k$ and $\sum_{i=1}^k p_i = 1$, i.e. the functions $\{p_i\}_{i=1, \dots, k}$ can be treated as place-dependent probabilities.

Moreover, we shall assume that there exists a set $X \subset \mathbb{R}^M$ such that

(a) X is a compact set with $\lambda_0 := (\int_X g_1, \dots, \int_X g_M) \in X$,

and for every $i = 1, \dots, k$:

(b) $F_i(X) \subset X$,

(c) $p_i|_X > 0$,

(d) $F_i|_X$ is a Lipschitz function with the Lipschitz constant $c_i < 1$.

Then the following assertions hold:

(1) $\mathcal{F} = (F_i, p_i)_{i=1}^k$ is an *iterated function system* on X .

(2) The IFS \mathcal{F} generates the following operator V acting on $M(X)$ (the space of all probability measures on X):

$$(V\nu)(B) = \sum_{i=1}^k \int_{F_i^{-1}(B)} p_i(\lambda) d\nu(\lambda) \quad (6.3)$$

for $\nu \in M(X)$ and $B \in B(X)$, where $B(X)$ denotes the family of all Borel sets on X . This operator describes the *evolution of probability measures* under the action of \mathcal{F} . We shall denote by $(Z_n^\nu)_{n \in \mathbb{N}}$ the associated Markov stochastic process having the initial distribution ν .

(3) There is a unique *invariant probability measure* μ for the IFS defined above fulfilling the equation $V\mu = \mu$. This measure is attractive, i.e. $V^n\nu$ converges weakly to μ for every $\nu \in M(X)$ as $n \rightarrow \infty$.

(4) The relative entropies G_n are given by

$$G_n = \int_X h_k(p_1(\lambda), \dots, p_k(\lambda)) d(V^n \delta_{\lambda_0})(\lambda) \quad \text{for } n \in \mathbb{N} \quad (6.4)$$

where h_k is the Shannon–Boltzmann entropy function given by $h_k(p_1, \dots, p_k) = -\sum_{i=1}^k p_i \ln p_i$ for any $p_i \geq 0$ such that $\sum_{i=1}^k p_i = 1$.

(5) The CS-entropy H_{meas} is given by an *integral formula*

$$H_{\text{meas}} = \int_X h_k(p_1(\lambda), \dots, p_k(\lambda)) \, d\mu(\lambda). \tag{6.5}$$

Let us sketch briefly the proof of the above statements. Assertion (1) follows from (0) and assumption (b). The Markov processes generated by IFSs were analysed in [64] and [67]. Assertion (3) can be deduced from assumptions (c) and (d), and [65, theorem 2.1]. Formulae (6.4) and (6.5) were proved by Fannes *et al* in [56] for algebraic measures, i.e. under the assumption that the formula for probabilities analogous to (2.19) holds. They followed an earlier result of Blackwell [68] on the entropy of functions of a finite-state Markov chain. In both these papers, however, the authors did not refer to the theory of IFSs and assumed that the matrices $D(i)$ are positive. In spite of this, their proof can also be applied in our case. For more details we refer the reader to [55].

6.2. Ergodic theorem and random algorithm for computing CS-entropy

The key point in our reasoning is to find a set X fulfilling conditions (a)–(d) above. In all the cases we analysed this task was not too difficult to accomplish. We shall give some examples below. Utilizing the results presented in [64] and [69] we can go even further and prove (under some additional assumptions) that G_n tends to H_{meas} exponentially. Moreover, applying the *Kajiser–Elton ergodic theorem for IFSs* (see [70] and [71]) we obtain the following formula:

$$H_{\text{meas}} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{l=0}^{n-1} h(Z_l^\nu) \quad \text{almost everywhere} \tag{6.6}$$

where $h = h_k(p_1, \dots, p_k)$ and ν is an arbitrary initial distribution.

This formula gives another numerical method of computing CS-entropy. To obtain the value H_{meas} it suffices to calculate Cesaro means of the function h along a trajectory of the stochastic process $(Z_l^\nu)_{l \in \mathbb{N}}$. This is a particular case of the general method which appeared under the name of random iterated algorithm in [66]. The convergence in (6.6) seems to be rather slow, namely as $n^{1/2}$. Note, however, that here the time computational complexity grows with k (the number of elements of the partition) linearly, whereas in the ‘matrix method’ we considered in sections 2.5 and 4.2 it grows polynomially (as k^n). Hence the method based on formula (6.6) may be specially useful for large values of k .

6.3. Example

Now let us consider the partition \mathcal{C}_3 of the sphere into three zones of equal volume: $E_1 = \{(\vartheta, \varphi) : \varphi \in [0, 2\pi), \vartheta \in [0, \pi/3]\}$, $E_2 = \{(\vartheta, \varphi) : \varphi \in [0, 2\pi), \vartheta \in (\pi/3, 2\pi/3]\}$, and $E_3 = \{(\vartheta, \varphi) : \varphi \in [0, 2\pi), \vartheta \in (2\pi/3, \pi]\}$. Set $j = \frac{1}{2}$. Then applying formula (4.3) one can show that the matrices $D(1), D(2), D(3)$ are given by

$$D(1) = \begin{pmatrix} \frac{1}{3} & \frac{2}{9} \\ \frac{2}{9} & \frac{13}{81} \end{pmatrix} \quad D(2) = \begin{pmatrix} \frac{1}{3} & 0 \\ 0 & \frac{1}{81} \end{pmatrix} \quad D(3) = \begin{pmatrix} \frac{1}{3} & -\frac{2}{9} \\ -\frac{2}{9} & \frac{13}{81} \end{pmatrix}. \tag{6.7}$$

Hence and from (6.1), (6.2) we obtain

$$p_1(\lambda) = \frac{1}{3} + \frac{2}{9}\lambda \quad p_2(\lambda) = \frac{1}{3} \quad p_3(\lambda) = \frac{1}{3} - \frac{2}{9}\lambda \tag{6.8}$$




Figure 9. The attracting invariant set for the IFS generated by the partition of the sphere into three zones of equal volume for (a) $j = \frac{1}{2}$, (b) $j = 1$. The values of the CS-measurement entropy computed by the random iterated algorithm are (a) $H_{\text{meas}} = 1.05306\dots$, (b) $H_{\text{meas}} = 0.99220\dots$

and

$$\begin{aligned}
 F_1(\lambda) &= (18 + 13\lambda)/(27 + 18\lambda) \\
 F_2(\lambda) &= \lambda/27 \\
 F_3(\lambda) &= (-18 + 13\lambda)/(27 - 18\lambda).
 \end{aligned} \tag{6.9}$$

The set $X = [-1, 1]$ fulfils conditions (a)–(d) with the contraction rates for the maps F_1 , F_2 , and F_3 equal to $c_1 = \frac{1}{3}$, $c_2 = \frac{1}{27}$, and $c_3 = \frac{1}{3}$, respectively. The support of the attracting invariant measure μ presented in figure 9(a) is a Cantor-like fractal set.

Now let us consider the case $j = 1$ (with the same partition \mathcal{C}_3). Applying formulae (4.3), (6.1), and (6.2), we can compute the maps p_1, p_2, p_3 , and F_1, F_2, F_3 , as before. Now, the set $X = \{(\lambda_1, \lambda_2) : \lambda_1 \in [-1, 1], \lambda_1^2 \leq \lambda_2 \leq 1\}$ satisfies conditions (a)–(d). The attractive invariant set for this IFS is presented in figure 9(b). Also in this case it has a fractal structure. The view of the middle part of the IFS through a magnifying glass is shown in figure 9(b) to underline its self-similar structure. In the figure caption we give the values of the CS-measurement entropy obtained with the aid of the random algorithm.

We have also applied this technique to compute numerically the CS-measurement entropy for other partitions of the phase space and j ranging from $\frac{1}{2}$ to 10. For the partition of the sphere into two hemispheres the results obtained in this way coincide with those received from the extrapolation of the relative entropies G_n and collected in table 1.

7. Rényi CS-measurement entropy

In this section we consider quantities which are natural generalizations of CS-measurement entropy introduced in section 2.3. We assume that \mathcal{C} is a partition of the phase space and

the CS-probabilities are given by (2.8). We shall write H_n for $H_n(\mathcal{C})$, G_n for $G_n(\mathcal{C})$, and H_{meas} for $H_{\text{meas}}(\mathcal{C})$. Moreover, we choose the parameter $\beta > 0$ such that $\beta \neq 1$.

There are at least two different ways of introducing a Rényi-type version of CS-measurement entropy. Firstly, we can define *CS-measurement entropy of order β* as

$$H_{\text{meas}}(\beta) := \limsup_{n \rightarrow \infty} \frac{1}{n} H_n(\beta) \tag{7.1}$$

where

$$H_n(\beta) := \frac{1}{1 - \beta} \ln \left[\sum_{i_0, \dots, i_{n-1}=1}^k (P_{i_0, \dots, i_{n-1}}^{\text{CS}})^\beta \right]. \tag{7.2}$$

On the other hand, using the notion of Rényi conditional entropy of order β [63] we can define the quantity

$$G_{\text{meas}}(\beta) := \limsup_{n \rightarrow \infty} G_n(\beta) \tag{7.3}$$

where

$$G_n(\beta) := \begin{cases} H_1(\beta) & \text{for } n = 1 \\ \frac{1}{1 - \beta} \ln \left[\sum_{i_0, \dots, i_{n-1}=1}^k (P_{i_0, \dots, i_{n-1}}^{\text{CS}})^\beta (P_{i_0, \dots, i_{n-2}}^{\text{CS}})^{1-\beta} \right] & \text{for } n > 1. \end{cases} \tag{7.4}$$

The quantities $G_n(\beta)$ are the analogues of the relative entropies considered in section 2. Note that $H_n(\beta) \rightarrow H_n(\beta \rightarrow 1)$ and $G_n(\beta) \rightarrow G_n(\beta \rightarrow 1)$. This justifies the notation $H_n(1) := H_n$, $G_n(1) := G_n$, and $H_{\text{meas}}(1) = G_{\text{meas}}(1) := H_{\text{meas}}$. In contrast to the case $\beta = 1$, the quantities $H_{\text{meas}}(\beta)$ and $G_{\text{meas}}(\beta)$ need not be equal in general (see figure 11).

The number $G_n(\beta)$ ($\beta \neq 1$) can be computed from the following integral formula analogous with (6.5):

$$G_{\text{meas}}(\beta) = \frac{1}{1 - \beta} \ln \int_X \sum_{i=1}^k (p_i(x))^\beta d\mu(x) \tag{7.5}$$

where $(X, (F_i)_{i=1}^k, (p_i)_{i=1}^k)$ is the iterated function system defined in section 6 and μ is the attractive invariant measure for this system [55].

Now let us consider the case of the division of the sphere into two hemispheres. As in section 4.4 we can use the Markovian approximation $G_2(\beta)$ to evaluate the limiting value $G_{\text{meas}}(\beta)$ for large values of j . Similar reasoning leads to the formula

$$G_2(\beta) = \frac{1}{1 - \beta} \ln \left[\frac{1 + \tau_j^\beta}{(1 + \tau_j)^\beta} \right] \tag{7.6}$$

where $\beta \neq 1$ and τ_j is given by formula (4.7).

The function G_2 defined by (4.8) and (7.6) is continuous. Moreover, we can compute the limits $G_2(\beta) \rightarrow \ln 2$ ($\beta \rightarrow 0$) and $G_2(\beta) \rightarrow \ln(1 + \tau_j)$ ($\beta \rightarrow \infty$). Asymptotically (for large j) we obtain

$$G_2(\beta) \sim \begin{cases} \frac{1}{1 - \beta} \frac{1}{(2\pi j)^{\beta/2}} & \text{for } \beta < 1 \\ \frac{\ln j}{2(2\pi j)^{1/2}} & \text{for } \beta = 1 \\ \frac{\beta}{\beta - 1} \frac{1}{(2\pi j)^{1/2}} & \text{for } \beta > 1. \end{cases} \tag{7.7}$$




Figure 10. Markovian approximation G_2 of the Rényi CS-measurement entropy as a function of the parameter β for the partition of the sphere into two hemispheres and selected values of the quantum number j labelling the curves.




Figure 11. Two versions of the Rényi CS-measurement entropy H_{meas} , G_{meas} , and the Markovian approximation G_2 as a function of the rescaled parameter $\zeta = 4 \arctan(\beta)/\pi$ for $j = 1$ and $j = 5$, in the case of the partition of the sphere into two hemispheres.

In figures 10 and 11 we treat the case of the partition of the sphere into two hemispheres. In figure 10 we present the Markovian approximation G_2 for different values of the semiclassical parameter j . We see that all the curves start from the value $\ln 2$ (topological entropy) and then decrease when the value of the parameter β grows. Moreover, we can observe that G_2 decreases when j increases and converges to 0 (which is the value of the classical Rényi entropy in this case) if j tends to ∞ . In figure 11 we compare two versions of the Rényi CS-measurement entropy $H_{\text{meas}}(\zeta)$, $G_{\text{meas}}(\zeta)$, and the Markovian approximation $G_2(\zeta)$ for two different values of the parameter j . The variable $\zeta = 4 \arctan(\beta)/\pi$ changes from 0 to 2, when β varies from 0 to ∞ . The quality of the Markovian approximation G_2 becomes worse for large values of β and j , still, it gives an upper bound for the Rényi CS-entropy G_{meas} .

8. Conclusions

This work has been devoted to the study of the notion of CS-measurement entropy. We have collected the basic theoretical material in sections 2.4 and 2.5, analysed numerical algorithms for computing CS-measurement entropy in sections 4.2 and 6, examined several examples in sections 4 and 5, and proposed two generalizations of the notion in section 7. The methods developed here can be used to investigate of the CS-measurement entropy for a broad class of partitions of the phase space and values of the semiclassical parameter j . The semiclassical limit (large j) has, as usual, been most difficult to treat. Nevertheless, even in this case, we have obtained several approximate results in section 4.4. We have restricted our attention to the spin ($SU(2)$) coherent states defined on the sphere S^2 . We believe, however, that our approach can be extended to other phase spaces and to other families of coherent states.

The fact that the measurement entropy H_{meas} can be calculated as the limit of the relative entropies G_n has played a crucial role in our analysis. As we have argued, the approach to the limit is exponential in this case. The rate of convergence seems to be strictly connected with the limiting value of the sequence: the larger the entropy H_{meas} , the faster the convergence. A similar dependence was reported for the KS-entropy of piecewise analytic one-dimensional maps by Szépfalusy and Györgyi [36]. They estimated the decay of the relative entropies G_n as $\sim e^{-2H(3)n}$, where $H(3)$ is the Rényi entropy of order 3. The convergence we have observed for CS-entropies is much faster.

In [13] we formulated a general programme for analysing *quantum chaos* in terms of CS-entropy. Here, we have studied CS-measurement entropy only, that is, the CS-entropy of the identity operator, which measures the randomness coming from the process of approximate sequential quantum measurement. Still, our main purpose is to study *CS-dynamical entropy*, which is connected only with the unitary dynamics of the quantum system and is defined as the difference of two quantities: the CS-entropy of the given unitary operator and the CS-measurement entropy (see formula (2.14)). The precise analysis of the notion of CS-measurement entropy is the first essential stage in performing this task. We expect that the methods elaborated here can also be used in the investigation of the CS-entropy for an arbitrary unitary map U , and so, in studying CS-dynamical entropy. The main difficulty in extending our approach to the general case is that we have to deal with much larger matrices; nevertheless, the numerical algorithms can be managed in much the same way. In a forthcoming publication we shall try to calculate the CS-dynamical entropy for quantized regular and chaotic maps.

In this work we have presented an effective method of computing the dynamical entropy of a system via *iterated function systems*. Although this technique has been applied here only in calculations of the CS-measurement entropy, we believe that it may be useful for computing the CS-dynamical entropy of quantum systems, as well as the Kolmogorov–Sinai entropy of classical systems.

Acknowledgments

We thank Mark Fannes, Piotr Garbaczewski, Denes Petz, Franklin Schroeck, and Tomasz Zastawniak for enlightening discussions. This work was supported by the Polish KBN grant 2 P03A 060 09.

Appendix A. Bounds for CS-measurement entropy

Proof of the inequalities (2.16)

We assume that \mathcal{C} denotes a finite partition of the phase space and we put $H_{\text{meas}} := H_{\text{meas}}(\mathcal{C})$, $H_n := H_n(\mathcal{C})$, and $G_n := G_n(\mathcal{C})$.

It follows from the general theory of dynamical entropy [50, 51] that the sequence $\{H_n\}_{n \in \mathbb{N}}$ is subadditive, i.e.

$$H_{n+l} \leq H_n + H_l \quad \text{for } n, l \in \mathbb{N}. \quad (\text{A1})$$

Let now $n, l \in \mathbb{N}$, $i_0, \dots, i_{n+l-1} = 1, \dots, k$. Then from (2.8) we deduce

$$\begin{aligned} P_{i_0, \dots, i_{n+l-1}}^{\text{CS}} &= \int_{E_{i_0}} dx_0 \dots \int_{E_{i_{n-1}}} dx_{n-1} \prod_{u=1}^{n-1} K(x_{u-1}, x_u) \int_{E_{i_n}} dx_n K(x_{n-1}, x_n) \\ &\quad \times \int_{E_{i_{n+1}}} dx_{n+1} \dots \int_{E_{i_{n+l-1}}} dx_{n+l-1} \prod_{u=n+1}^{n+l-1} K(x_{u-1}, x_u) \\ &\leq P_{i_0, \dots, i_{n-1}}^{\text{CS}} K_0 P_{i_n, \dots, i_{n+l-1}}^{\text{CS}}. \end{aligned} \quad (\text{A2})$$

Taking the logarithms of both sides of (A2), multiplying them by $-P_{i_0, \dots, i_{n+l-1}}^{\text{CS}}$, and summing over $i_0, \dots, i_{n+l-1} = 1, \dots, k$ we get

$$H_{n+l} \geq H_n - \ln K_0 + H_l \quad \text{for } n, l \in \mathbb{N}. \quad (\text{A3})$$

Combining (A1) with (A3) and dividing the expressions by n we have

$$\frac{1}{n} H_n - \frac{1}{n} \ln K_0 \leq \frac{1}{n} (H_{l+n} - H_l) \leq \frac{1}{n} H_n \quad (\text{A4})$$

and so

$$\frac{1}{n} H_n - \frac{1}{n} \ln K_0 \leq \frac{1}{n} \sum_{i=1}^n G_{l+i} \leq \frac{1}{n} H_n. \quad (\text{A5})$$

Letting $l \rightarrow \infty$ we obtain the desired conclusion.

Appendix B. Convergence rate of partial entropies

Proof of property 2.4 (4)

We follow the notation of appendix A. From 2.4 (1) we get

$$\frac{1}{n} H_n = \frac{1}{n} \sum_{i=1}^n G_i \geq \frac{1}{n} H_1 + \frac{n-1}{n} H_{\text{meas}}. \quad (\text{B1})$$

Hence

$$\frac{1}{n} H_n - H_{\text{meas}} \geq \frac{H_1 - H_{\text{meas}}}{n} > 0. \quad (\text{B2})$$

On the other hand (2.16) implies

$$\frac{1}{n} H_n - H_{\text{meas}} \leq \frac{\ln K_0}{n}. \quad (\text{B3})$$

Combining (B1) and (B2) we get the required result.

Appendix C. Formula for the second-order CS-probabilities

Proof of formula (4.6).

Set $2j = M$. Then, from (4.2) and (4.3) we have

$$\begin{aligned}
 P_{+-}^{\text{CS}} &= \int_0^1 \frac{dt}{2} \int_{-1}^0 \frac{ds}{2} \tilde{K}(t, s) \\
 &= \frac{M+1}{4^{M+1}} \sum_{q=0}^M \binom{M}{q}^2 \int_0^1 dt (1+t)^q (1-t)^{M-q} \int_{-1}^0 ds (1+s)^q (1-s)^{M-q} \\
 &= \frac{1}{(M+1)4^{M+1}} \sum_{q=0}^M R_q^M R_{M-q}^M
 \end{aligned} \tag{C1}$$

where

$$R_p^M := (M+1) \binom{M}{p} \int_0^1 dt (1+t)^p (1-t)^{M-p}. \tag{C2}$$

Now we need the following two lemmas, which we shall prove later.

Lemma 1.

$$R_p^M = \sum_{s=0}^p \binom{M+1}{s} \tag{C3}$$

and

Lemma 2.

$$\sum_{q=0}^M R_q^M R_{M-q}^M = (2M+1) \binom{2M}{M}. \tag{C4}$$

Combining (C1) and (C4) we get

$$P_{+-}^{\text{CS}} = \frac{1}{(M+1)4^{M+1}} (2M+1) \binom{2M}{M} = \binom{2M+1}{M} \frac{1}{4^{M+1}} \tag{C5}$$

which establishes the formula.

Proof of lemma 1. We proceed by induction. Clearly, $R_0^M = 1$. Assuming (C3) to hold for p , we shall prove it for $p+1$. We have

$$R_{p+1}^M = (M+1) \binom{M}{p+1} \int_0^1 dt (1+t)^{p+1} (1-t)^{M-p-1}. \tag{C6}$$

Integrating by parts we obtain

$$R_{p+1}^M = \frac{M+1}{M-p} \binom{M}{p+1} + \frac{(M+1)(p+1)}{M-p} \binom{M}{p+1} \int_0^1 dt (1+t)^p (1-t)^{M-p}. \tag{C7}$$

By the induction assumption

$$R_{p+1}^M = \binom{M+1}{p+1} + R_p^M \tag{C8}$$

which completes the proof.

Proof of lemma 2. Applying lemma 1 we deduce that

$$\begin{aligned} \sum_{q=0}^M R_q^M R_{M-q}^M &= \sum_{\substack{s,l=0; \\ s+l \leq M}}^M \binom{M+1}{s} \binom{M+1}{l} ((M+1) - (s+l)) \\ &= \sum_{r=0}^{M+1} \sum_{s=0}^r \binom{M+1}{s} \binom{M+1}{r-s} ((M+1) - r). \end{aligned} \quad (\text{C9})$$

Using the well known combinatorial identities

$$\sum_{s=0}^L \binom{L}{s} \binom{L}{r-s} = \binom{2L}{r} \quad (\text{C10})$$

$$\sum_{r=0}^L \binom{2L}{r} = \frac{1}{2} \left(4^L + \binom{2L}{L} \right) \quad (\text{C11})$$

$$\sum_{r=0}^L \binom{2L}{r} r = \frac{L}{2} 4^L \quad (\text{C12})$$

we conclude that

$$\begin{aligned} \sum_{q=0}^M R_q^M R_{M-q}^M &= \sum_{r=0}^{M+1} \binom{2M+2}{r} ((M+1) - r) \\ &= \frac{M+1}{2} \binom{2M+2}{M+1} = (2M+1) \binom{2M}{M} \end{aligned} \quad (\text{C13})$$

which proves the lemma.

References

- [1] Balazs N L and Voros A 1989 The quantized Baker's transformation *Ann. Phys., NY* **190** 1
- [2] Saraceno M 1990 Classical structures in the quantized Baker transformation *Ann. Phys., NY* **199** 37
- [3] De Bièvre S, Degli Epositi M and Giachetti R 1996 Quantization of a class of piecewise affine transformations on the torus *Commun. Math. Phys.* **176** 73
- [4] Hannay J H and Berry M V 1980 Quantization of linear maps on a torus—Fresnel diffraction by a periodic grating *Physica* **1D** 267
- [5] Haake F, Kuś M and Scharf R 1987 Classical and quantum chaos for a kicked top *Z. Phys. B* **65** 381
- [6] Frahm H and Mikeska H J 1986 Level statistics and stochasticity in a driven quantum system *Z. Phys. B* **65** 249
- [7] Nakamura K, Okazaki Y and Bishop A R 1986 Periodically pulsed spin dynamics: scaling behaviour of semiclassical wavefunctions *Phys. Rev. Lett.* **57** 5
- [8] Haake F 1991 *Quantum Signatures of Chaos* (Berlin: Springer)
- [9] Gutzwiller M C 1991 *Chaos in Classical and Quantum Mechanics* (Berlin: Springer)
- [10] Peres A 1993 *Quantum Theory: Concepts and Methods* (Dordrecht: Kluwer)
- [11] Wehrl A 1991 The many facets of entropy *Rep. Math. Phys.* **30** 119
- [12] Ohya M and Petz D 1993 *Quantum Entropy and Its Use* (Berlin: Springer)
- [13] Stomczyński W and Życzkowski K 1994 Quantum chaos: an entropy approach *J. Math. Phys.* **35** 5674
- [14] Roepstorff G 1995 Quantum dynamical entropy *Chaos. The Interplay Between Stochastic and Deterministic Behaviour, Proc. XXXIst Winter School of Theoretical Physics (Karpacz, 13–24 February)* ed P Garbaczewski et al (Berlin: Springer) pp 305–12
- [15] Vilela Mendes R 1995 Entropy and quantum characteristic exponents *Chaos. The Interplay Between Stochastic and Deterministic Behaviour, Proc. XXXIst Winter School of Theoretical Physics (Karpacz, 13–24 February)* ed P Garbaczewski et al (Berlin: Springer) pp 273–82
- [16] Connes A, Narnhoffer H and Thirring W 1987 Dynamical entropy of C^* -algebras and von Neumann algebras *Commun. Math. Phys.* **112** 691

- [17] Gaspard P 1994 Comment on dynamical randomness in quantum systems *Prog. Theor. Phys. Suppl.* **116** 369
- [18] Alicki R and Fannes M 1994 Defining quantum dynamical entropy *Lett. Math. Phys.* **32** 75
- [19] Roepstorff G 1995 Quantum dynamical entropy based on histories *Preprint RWTH Aachen, Aachen*
- [20] Davies E B 1976 *Quantum Theory of Open Systems* (London: Academic)
- [21] Prugovečki E 1986 *Stochastic Quantum Mechanics and Quantum Spacetime* (Dordrecht: Reidel)
- [22] Busch P, Lahti P J and Mittelstaedt P 1991 *The Quantum Theory of Measurement* (Berlin: Springer)
- [23] Schroeck F E Jr 1995 *Quantum Mechanics on Phase Space* (Dordrecht: Kluwer)
- [24] Lüders G 1951 Über die zustandsänderung durch den meßprozeß *Ann. Phys., Lpz.* **8** 322
- [25] Ozawa M 1985 Conditional probability and *a posteriori* states in quantum mechanics *Publ. RIMS, Kyoto Univ.* **21** 279
- [26] Schroeck F E Jr 1982 On the stochastic measurement of incompatible spin components *Found. Phys.* **12** 479
- [27] Busch P 1986 Unsharp reality and joint measurements for spin observables *Phys. Rev. D* **33** 2253
- [28] Busch P and Schroeck F E Jr 1989 On the reality of spin and helicity *Found. Phys.* **19** 807
- [29] Garbaczewski P 1990 Randomness in the quantum description of neutral spin 1/2 particles *Fortschr. Phys.* **38** 447
- [30] Eckmann J-P and Ruelle D 1985 Ergodic theory of chaos and strange attractors *Rev. Mod. Phys.* **57** 617
- [31] Christiansen F and Politi A 1995 Generating partition for the standard map *Phys. Rev. E* **51** R3811
- [32] Życzkowski K and Słomczyński W 1995 Exponential decay of relative entropies to the Kolmogorov–Sinai entropy for the standard map *Phys. Rev. E* **52** 6879
- [33] Mañé R 1987 *Ergodic Theory and Differentiable Dynamics* (Berlin: Springer)
- [34] Misiurewicz M and Ziemian K 1987 Rate of convergence for computing entropy of some one-dimensional maps, *Proc. Conf. Ergodic Theory and Related Topics II (Georgenthal, April 20–25, 1986)* ed H Michel (Leipzig: BSB Teubner) pp 147–59
- [35] Ziemian K 1989 Rate of convergence of conditional entropies for some maps of an interval *Stud. Math.* **93** 259
- [36] Szépfalussy P and Györgyi G 1986 Entropy decay as a measure of stochasticity in chaotic systems *Phys. Rev. A* **33** 2852
- [37] Szépfalussy P 1989 Characterization of chaos and complexity by properties of dynamical entropies *Phys. Scr. T* **25** 226
- [38] Csordás A, Györgyi G, Szépfalussy P and Tél T 1993 Statistical properties of chaos demonstrated in a class of one-dimensional maps *Chaos* **3** 31
- [39] Ebeling W, Freund J and Rateitschak K 1996 Entropy and extended memory in discrete chaotic dynamics *Int. J. Bifurc. Chaos* in press
- [40] Rateitschak K, Freund J and Ebeling W 1996 Entropy of sequences generated by nonlinear processes: The logistic map *Entropy and Entropy Generation: Fundamentals and Applications* ed J S Shiner (Dordrecht: Kluwer)
- [41] Cohen A and Procaccia I 1985 Computing the Kolmogorov entropy from time signals of dissipative and conservative dynamical systems *Phys. Rev. A* **31** 1872
- [42] Klauder J R and Skagerstam B 1985 *Coherent States* (Singapore: World Scientific)
- [43] Davies E B and Lewis J T 1970 An operational approach to quantum probability *Commun. Math. Phys.* **17** 239
- [44] Ozawa M 1993 Canonical approximate quantum measurements *J. Math. Phys.* **34** 5596
- [45] Aczél J 1984 Measuring information beyond communication theory—Why some generalized information measures may be useful, others not *Aequationes Math.* **27** 1
- [46] Kapur J N 1994 *Measures of Information and Their Applications* (New York: Wiley)
- [47] Weaire D and Williams A R 1977 The Anderson localization problem: I. A new numerical approach *J. Phys. C: Solid State Phys.* **10** 1239
- [48] Życzkowski K and Słomczyński W 1995 Coherent states quantum entropy *Proc. Int. Conf. on Dynamical Systems and Chaos (Tokyo, 23–27 May, 1994)* vol 2, ed Y Aizawa *et al* (Singapore: World Scientific) pp 467–70
- [49] Słomczyński W 1997 From quantum entropy to iterated function systems *Chaos, Solitons & Fractals* in press
- [50] Martin N F G and England J W 1981 *Mathematical Theory of Entropy* (Reading, MA: Addison-Wesley)
- [51] Kifer Y 1988 *Random Perturbations of Dynamical Systems* (Boston, MA: Birkhäuser)
- [52] Perelomov A 1986 *Generalized Coherent States and Their Applications* (Berlin: Springer)
- [53] Zhang W-M, Feng D H and Gilmore R 1990 Coherent states: theory and some applications *Rev. Mod. Phys.* **62** 867
- [54] Zhang W-M and Feng D H 1995 Quantum nonintegrability in finite systems *Phys. Rep.* **252** 1
- [55] Słomczyński W Dynamical entropy and iterated function systems, in preparation

- [56] Fannes M, Nachtergaele B and Slegers L 1992 Functions of Markov processes and algebraic measures *Rev. Mod. Phys.* **4** 39
- [57] Radcliffe J M 1971 Some properties of coherent spin states *J. Phys. A: Math. Gen.* **4** 313
- [58] Arecchi F T, Courtens E, Gilmore R and Thomas H 1972 Atomic coherent states in quantum optics *Phys. Rev. A* **6** 2211
- [59] Hecht K T 1987 *The Vector Coherent State Method and Its Application to Problems of Higher Symmetry (Lecture Notes in Physics 290)* (Berlin: Springer)
- [60] Vieira V R and Sacramento P D 1995 Generalized phase-space representatives of spin- J operators in terms of Bloch coherent states *Ann. Phys., NY* **242** 188
- [61] Schroeck F E Jr 1985 The dequantization programme for stochastic quantum mechanics *J. Math. Phys.* **26** 306
- [62] Werner R F and Wolff M P H 1995 Classical mechanics as quantum mechanics with infinitesimal \hbar *Phys. Lett. A* **202** 155–9
- [63] Rényi A 1961 On measures of entropy and information *Proc. 4th Berkeley Symp. Math. Statist. Prob. (1960)* vol I (Berkeley, CA: University of California Press) pp 547–61
- [64] Iosifescu M and Grigorescu G 1990 *Dependence with Complete Connections and its Applications* (Cambridge: Cambridge University Press)
- [65] Barnsley M F 1988 *Fractals Everywhere* (Boston, MA: Academic)
- [66] Barnsley M F 1989 Iterated function systems *Chaos and Fractals. The Mathematics Behind the Computer Graphics* ed R L Devaney and L Keen (Providence, RI: American Mathematical Society) pp 127–44
- [67] Barnsley M F, Demko S G, Elton J H and Geronimo J S 1988 Invariant measures for Markov processes arising from iterated function systems with place-dependent probabilities *Ann. Inst. H Poincaré* **24** 367
- [68] Blackwell D 1957 The entropy functions of finite-state Markov chains *Trans. 1st Prague Conf. Information Theory, Statistical Decision Functions, Random Processes (Liblice, November 28–30, 1956)* (Prague: Czechoslovak Academy of Sciences) pp 13–20
- [69] Lasota A and Yorke J A 1994 Lower bound technique for Markov operators and iterated function systems *Random Comput. Dynam.* **2** 41
- [70] Kaijser T 1981 On a new contraction condition for random systems with complete connections *Rev. Roum. Math. Pure Appl.* **26** 1075
- [71] Elton J H 1987 An ergodic theorem for iterated maps *Ergod. Theor. Dynam. Sys.* **7** 481